

Hints of $\mu + N \rightarrow \mu + N + \mu^+ \mu^-$ process observation with the MACRO Experiment

G. Battiston^{i*} and E. Scapparone⁺ for the MACRO Collaboration

^{*}INFN Milano, via Celoria 16, 20133, Italy

⁺ INFN - Laboratori Nazionali del Gran Sasso, S.S.17 km 18+910, 61070, Assergi (AQ), Italy

**Talk at Xth International Symposium on Very High
Energy Cosmic Ray Interaction
July 12 - 17, 1998
To be published in Nucl. Phys. B, Proc. Suppl.**

Hints of $\mu+N \rightarrow \mu+N+\mu^+\mu^-$ process observation with the MACRO Experiment

G. Battistoni^a and E. Scapparone^b for the MACRO Collaboration *

^aINFN Milano, via Celoria, Italy

^bINFN, LNGS S.S.17 km 18+910, 61070, Assergi (AQ), Italy

We present the analysis of the distribution of the distance of muon pairs detected deep underground by the MACRO experiment. A detailed analysis of the low separation region pointed out an excess of experimental data with respect to the Monte Carlo expectation. The effect is discussed taking into account the $\mu+N \rightarrow \mu+N+\mu^+\mu^-$ process.

1. Introduction

The distribution of the muon pair separation deep underground (decoherence function) plays an important role in the study of Cosmic Ray physics in underground experiments, being sensitive to some aspect of the hadronic interaction model. Taking advantage of the large area, MACRO[1] can access the largest separation, exploring thus the highest P_t regions. Moreover, the capability of the MACRO detector to resolve very close tracks permits the extension of the decoherence analysis to a distance region hardly studied in the past.

2. Analysis of the behaviour at small distances

We have analyzed about $3.4 \cdot 10^5$ events, corresponding to 7732 hr live time using the lower part of the apparatus. These events were submitted to the following selection criteria:

- Zenith angle smaller than 60° : this choice is dictated by our limited knowledge of the Gran Sasso topographical map for high zenith angles. Furthermore, we cannot disregard the atmosphere curvature for larger zenith angles, while at present our simulation model does not include

that.

- The number of streamer tube hits out of track smaller than 45. This selection aims to eliminate possible misleading track reconstruction in events produced by noise in the streamer tube system and/or e.m. interactions inside or in the neighbourhood of the apparatus.
- Track pairs which survive a loose parallelism cut: this rejects hadrons from photonuclear interactions close to the detector. This selection also eliminates tracks reconstructed using hits due to e.m. interactions in events survived to the previous cut.

The last cut is not completely efficient in rejecting muon tracks originating from local particle production because the angle between these tracks may fall within the limits imposed by the parallelism cut. Moreover, these limits cannot be further reduced since the average angular divergence due to multiple muon scattering in the rock overburden, is about 1° at the MACRO depth.

A further selection was applied in order to reduce these effects. We computed, for each muon track in the wire view, the ratio R between the number of streamer tube planes fired by the muon to the number of expected fired planes considering the track direction. Only tracks with $R \geq 0.75$ were accepted. The application of this cut (hereafter the C4 cut) in the wire view alone is a good compromise between the rejection capability of the algorithm and the loss of events due to

*For the complete list of the Collaboration see the paper "Relevance of the hadronic interaction model in the interpretation of multiple muon data as detected with MACRO experiment" by O. Palamara at these proceedings




Figure 1. *Comparison between the experimental decoherence function with the HEMAS expectation.*

the unavoidable inefficiency of the streamer tube system. We found that in the wire projective view the probability to reject a muon track for a casual alignment contiguous inefficient planes is 2.0%.

To test the ability of this cut to reject hadronic tracks, we used FLUKA [5] to simulate 3028 hr of live time in which muons were accompanied by hadronic products of photonuclear interactions in the 10 m of rock surrounding the detector. We found that the parallelism cut alone provides a rejection efficiency of about 54.6% of the pair sample, while the addition of cut C4 enhances the rejection to 95.9%. Furthermore the effect of hadron contamination, is very small, contributing less than 1% in the overall muon pair sample. After the overall application of these cuts, the number of unambiguously associated muon pair tracks survived is 355,795. Fig. 1 shows the comparison between real data and HEMAS expectation: a general good agreement is found, showing the HEMAS capability in reproducing the muon bundle features deep underground. Nevertheless a relevant discrepancy at low distances is observed, indicating a possible physical source for this excess in the real data. For instance, considering separation smaller than 80 cm, an excess of 55% is observed in the real data (Table 1). The pro-

cess of muon pair production by muons in the rock is a natural candidate. As pointed out in [4], at the typical muon energy involved in underground analyses ($E_\mu \sim 1$ TeV), if very large energy transfer are considered, the cross section of such a process is non negligible with respect to the e^+e^- pair production. The key point is that at high $v=E_\gamma/E_\mu$, the most important Feynmann diagram is rather insensitive to the ratio $(m_e/m_\mu)^2$. An analytic expression for the muon pair production cross section is given in [4,8]. In order to test the hypothesis, such cross section has been included in the muon transport code PROPMU[7]. Assuming a muon flux with energy spectrum $E^{-3.7}$ with a minimum muon energy $E_\mu^{min} = 1.2$ TeV at the surface and considering the actual mountain profile, we generated a sample of 10^7 muons corresponding to 3666 h of live time. About $\sim 3.0 \cdot 10^6$ muons survived to the MACRO level, 5360 of which derived from muon pair production processes. The average distance of these muon pairs is (128 ± 1) cm and their average energies are (657 ± 14) GeV and (145 ± 3) GeV for, the main muon and the secondary muon samples respectively. We propagated the muons surviving to the MACRO level through GEANT simulation and we applied to these events the same cuts specified in Section 2. Finally, the number of events was normalized to the live time of real data.

In Table 1 we report the number of weighted muon pairs in the first bins of the experimental and simulated decoherence distributions expressed as percentage with respect to the population in the bin of the distribution maximum. The effect of standard cuts, of the C4 cut and of the subtraction of the muon pair production process are shown in order. In each case, we indicate in percentage the bin populations with respect to the peak of the distribution and the discrepancy with respect to the Monte Carlo predictions.

In Fig. 2 we compare the simulated decoherence curve with the experimental data corrected for the muon pair production effect. Despite the approximation introduced in our test, it seems that the proposed muon pair production process can account for the most of observed discrepancy in the low distance range.

Table 1

Number of weighted muon pairs in the first bins of the experimental and simulated decoherence distributions. The discrepancy is the percentage difference between experimental and Monte Carlo values normalized to the distribution maximum (last column).

	0–80 cm	80–160 cm	160–240 cm	240–320 cm	320–400 cm (max)
Exp. Data	5528	12491	17569	20514	20816
MC Data	5154	21417	33573	40367	42679
Discrepancy	(55±2)%	(16±2)%	(6±1)%	(4±1)%	
Exp. Data + C4	3612	11128	16535	19597	19977
MC Data + C4	4848	20346	31932	38425	40660
Discrepancy	(34±2)%	(10±2)%	(6±2)%	(4±2)%	
Exp. Data + C4 μ pair subtraction	2193	9264	15462	19190	19842
Discrepancy	(8±7)%	(7±3)%	(0±2)%	(2±2)%	




Figure 2. The low distance region of the experimental decoherence function before and after the subtraction of the secondary muon sample and comparison with the Monte Carlo simulation.

3. Discussion and conclusion

We measured the underground decoherence function using high energy muons ($E_\mu > 1.3$ TeV) up to a maximum distance of about 70 m.

The capability to resolve very close muon

tracks permitted the investigation of the behaviour of the decoherence function for small separations. Apart from the negligible contamination of hadro-production by muons, we found strong hints of a contribution coming from muon pair production by muons. The inclusion of this process in the simulation reproduces, both in a qualitative and quantitative way, the experimental data.

REFERENCES

- [1] MACRO Collaboration, S.P. Ahlen et al., Nucl. Instr. Meth. **A324** (1993) 337.
- [2] MACRO Collaboration, S.P. Ahlen et al., Phys. Rev. **D46** (1992) 4836.
- [3] MACRO Collaboration, M. Ambrosio et al., Proc. of the XXV Int. Cosmic Ray Conf., Durban, South Africa, 1997, Vol. 6, p. 357.
- [4] V.A. Kudryavtsev and O.G. Ryazhskaya, INFN-AE-97-08 (1997)
- [5] A. Fassó et al., An update about FLUKA, Proc. 2nd workshop on Simulating Accelerator Radiation Environment, SARE-2, CERN-Geneva, 9-11, October, 1995.
- [6] R. Brun et al., CERN GEANT3 User's guide, DD/EE/84-1 (1992).
- [7] P. Lipari and T. Stanev, Phys. Rev **D44** (1991) 3543.
- [8] S.R. Kelner, Sov. J. Nucl. Phys. **5** (1967) 778.